## Recapitulating pancreatic cancer ecosystems by multicellular cancer organoid cultures based on patient-derived cancer cells

Cancer ecosystems comprise not only of cancer cells but also of various stromal cells, such as mesenchymal and vascular endothelial cells. Pancreatic cancer is stromal-rich cancer, and the abundant stroma is considered to responsible for the resistance to treatment and for poor prognosis. Contrary to expectations, however, treatment targeting stromal mesenchymal cells exacerbates pancreatic cancer. Accordingly, it is important to understand and control pancreatic cancer cell- stromal cell interaction rather than merely targeting the stroma. In order to analyze the cancer cell- stromal cell interaction and further accurately assess the susceptibility of pancreatic cancer drugs, a culture system capable of reproducing pancreatic cancer ecosystem is essential.

We previously established a method to generate multi-lineage functional organs using human iPS cells. Autonomous cell aggregation is induced by co-culturing human iPS cell-derived hepatic endoderm, vascular endothelial cells, mesenchymal cells under certain conditions results in the generation of the liver primordia i.e. iPSC liver bud (Takebe T. Sekine K et al. Nature, 2013; Takebe T, Sekine K et al., Cell Reports, 2017). Furthermore, single-cell RNA sequencing analysis elucidated the dynamic cell-cell interaction of different cell types (Sekine K, Camp JG et al. Nature 2017).

By applying the three-dimensional tissue reconstruction technique to pancreatic cancer research, we succeeded in reconstituting human pancreatic cancer tissue (pancreatic cancer organoid) recapitulating abundant stroma from patient-derived primary cancer cells. Evaluation of drug susceptibility of pancreatic cancer organoids in vitro revealed that drug sensitivity to gemcitabine, a therapeutic drug for pancreatic cancer, is greatly reduced compared to the cancer cell. Transplantation of cancer organoid into immunodeficient mice generated pancreatic cancer xenografts with abundant stroma and pancreatic ductal structures, which resembled patient pancreatic cancer tissue. Furthermore, drug sensitivity of the xenograft was also significantly decreased. We found cancer cell- stromal cell interaction is responsible for the treatment resistance of pancreatic cancer organoids xenograft.

In summary, pancreatic cancer organoids are an effective tool for reproducing the treatment resistance of pancreatic cancer patients and useful for drug screening.

## [References]

- Takebe T\*, Treutlein B\* † equal contribution., \* corresponding author. Multilineage communication regulates human liver bud development from pluripotency Nature 546:533-538. (2017)
- 2. Takebe T, Sekine K, Enomura M, Koike H, Kimura M, Ogaeri T, Zhang RR, Ueno Y, Zheng YW, Koike N, Aoyama S, Adachi Y, Taniguchi H Vascularized and functional human liver tissue from an induced pluripotent stem cell-derived organ bud transplant.
  - Nature 499(7459):481-484. (2013)
- 3. Takahashi Y, Sekine K, Kin T, Takebe T, Taniguchi H Transplantation
  - Cell Reports 23(6):1620-1629. (2018)
- 4. Takebe T, Sekine K, Kimura M, Yoshizawa E, Ayano S, Koido M, Funayama S, Nakanishi N, Hisai T, Kobayashi T, Kasai T, Kitada R, Mori A, Ayabe H, Ejiri Y, Amimoto N, Yamazaki Y, Ogawa S, Ishikawa M, Kiyota Y, Sato Y, Nozawa K, Okamoto S, Ueno Y, Taniguchi H. Massive and reproducible production of liver buds entirely from human pluripotent stem cells Cell Reports 21(10):2661-2670. (2017)



## Keisuke SEKINE, PhD

Associate Professor, The University of Tokyo

2000 Ph.D., The University of Tokyo 2000 Research Associate, The University of Tokyo 2009 Assistant Professor, Yokohama City University 2018 Junior Associate Professor, Yokohama City University 2019 Associate Professor, The University of Tokyo



## Keisuke Sekine

Division of Regenerative Medicine, Center for Stem Cell Biology and Regenerative Medicine, The Institute of Medical Science, The University of Tokyo

1. Sekine K<sup>†</sup>, Camp JG <sup>†</sup>, Gerber T, Loeffler-Wirth H, Binder H, Gac M, Kanton S, Kageyama J, Damm G, Seehofer D, Belicova L, Bickle M, Barsacchi R, Okuda R, Yoshizawa E, Kimura M, Ayabe H, Taniguchi H,

Self-Condensation Culture Enables Vascularization Of Tissue Fragments For Efficient Therapeutic